

Durham Research Online

Deposited in DRO:

20 January 2016

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Liu, Jean and Milne, Glenn A. and Kopp, Robert E. and Clark, Peter U. and Shennan, Ian (2016) 'Sea-level constraints on the amplitude and source distribution of Meltwater Pulse 1A.', *Nature geoscience.*, 9 (2). pp. 130-134.

Further information on publisher's website:

<http://dx.doi.org/10.1038/ngeo2616>

Publisher's copyright statement:

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Sea-Level Constraints on the Amplitude and Source Distribution of Meltwater Pulse

1A

Jean Liu¹, Glenn A. Milne^{1*}, Robert E. Kopp², Peter U. Clark³, and Ian Shennan⁴

¹Department of Earth and Environmental Sciences, University of Ottawa

²Department of Earth & Planetary Sciences, Rutgers Energy Institute, and Institute of Earth,
Ocean & Atmospheric Sciences, Rutgers University

³ College of Earth, Ocean, and Atmospheric Sciences, Oregon State University

⁴Department of Geography, Durham University

*To whom correspondence should be addressed, gamilne@uottawa.ca

During the last deglaciation, sea levels rose as ice sheets retreated. This climate transition was punctuated by periods of more intense melting; the largest and most rapid of these – Meltwater Pulse 1A – occurred about 14,500 years ago, with rates of sea level rise reaching approximately 4 m per century¹⁻³. Such rates of rise suggest ice-sheet instability, but the meltwater sources are poorly constrained, thus limiting our understanding of the causes and impacts of the event⁴⁻⁷. In particular, geophysical modelling studies constrained by tropical sea-level records^{1,8,9} suggest an Antarctic contribution of more than seven meters, whereas most reconstructions from Antarctica indicate no substantial change in ice-sheet volume around the time of Meltwater Pulse 1A¹⁰. Here we use a glacial isostatic adjustment model to reinterpret tropical sea level reconstructions from Barbados², the Sunda Shelf³ and Tahiti¹. According to our results, global mean sea level rise during Meltwater Pulse 1A was between 8.6 and 14.6 metres (95% probability). As for the melt partitioning, we find an allowable contribution from Antarctica of either 4.1 to 10.0 m or 0 to 6.9 m (95% probability), using two recent

1 estimates^{11,12} of the contribution from the North American ice sheets. We conclude that a
2 significant contribution of melt from the Antarctic ice sheets is not necessarily required to
3 explain the documented sea level rise during Meltwater Pulse 1A.

4 Using a glacial isostatic adjustment (GIA) sea-level model (see Methods), global sea-
5 level changes for a wide range of ice histories were calculated and then compared to palaeo-sea
6 level reconstructions (based on sedimentary and geomorphological indicators, such as corals and
7 mangroves, including their uncertainties) to assess whether a given Meltwater Pulse 1A (MWP-
8 1A) source scenario is compatible with the field constraints. We focused on modelling the
9 relative sea level (rsl) change across MWP-1A in order to reduce the sensitivity of the analysis to
10 mantle viscosity structure, which is not precisely known. A primary limitation of this approach
11 is that there are only three far-field sites (locations in low latitudes distant from ice sheets) where
12 sea-level records constrain the amplitude of MWP-1A: Barbados, Sunda Shelf, and Tahiti.
13 Within this limited data framework, a key aim of this study is to quantify the possible MWP-1A
14 source constraints via a sea-level fingerprinting^{8,13} analysis when both data and model
15 uncertainty are taken into consideration.

16 MWP-1A was first identified at Barbados from reef framework-forming corals with
17 species-dependent depth ranges^{14,15,16}. By assuming that the coral growth could keep pace with
18 sea level during periods of rapid sea-level rise, previous work² estimated that MWP-1A occurred
19 between 14.2 ka and 13.5 ka and had a rsl amplitude of 14 -24 m. However, dated samples of the
20 shallow-water coral species (*Acropora palmata*) prior to 14.2 ka suggest that this interpretation
21 may be incorrect: specifically, the rate of sea-level rise had already increased prior to 14.2 ka
22 (ref. 11) and the shallow-water corals were already in the process of drowning due to rapid rates
23 of sea-level rise^{7,17}. A recent study of coral records from Tahiti supports the latter interpretation

by constraining MWP-1A to have occurred within the period 14.65-14.31 ka (ref. 1), thus defining a maximum duration of 340 years for the event. Adopting 14.31 ka as the end of MWP-1A, Deschamps et al.¹ estimated the Barbados MWP-1A amplitude to be ~15 m (Fig. 1). In this study, we extended this reappraisal by considering the depth and age uncertainties of the Barbados coral record, and arrived at a MWP-1A amplitude range of 9.7-33.6 m at this location (Fig. 1 and caption). The maximum estimate is large and could likely be reduced via a more sophisticated approach that considers additional information, such as reef morphology and stratigraphy; however, this is beyond the scope of this study. Furthermore, we note that the upper bound at this site does not play an important role in our final results (see below).

The Sunda Shelf record³ is defined by rooted mangrove trees, which, like corals, grow in a specific elevation range relative to mean sea level. Fossil mangrove roots were recovered from sediment cores distributed over a relatively large area of the shelf (Fig. 2a). Assuming that all the core sites reflect the same sea-level history suggests that MWP-1A had a rsl amplitude of 12-20 m and occurred at the same time as it did in Tahiti^{1,3} (Fig. 2b). There is, however, a considerable sea-level gradient across the region due to water loading associated with flooding of the shelf¹⁸ (Fig. 2a); this gradient influences the estimated MWP-1A amplitude since the core locations are widely separated. Using the GIA sea-level model introduced above with two alternative ice models and 162 combinations of earth-model parameters, we translated the Sunda Shelf observations to their equivalent values at a single location, site 18300 (Fig. 2a, black star; see Methods). The model results indicate that this translation leads to a 2 to 4 m correction in rsl at sites where samples define the beginning (Fig. 2a, blue dot) and end (Fig. 2a, red dot) of MWP-1A. Correcting for the spatial sea-level gradient yields a MWP-1A amplitude of 7.5 to 17.3 m (Fig. 2c), which is significantly reduced compared to the original interpretation³ (Fig. 2b).

1 The final observations we consider are from Tahiti, which, as described earlier, are from
2 a well-dated high-resolution coral record¹. A large number of cores were drilled, resulting in a
3 local sea-level record that agrees with a heterogeneous reef-accretion model²⁰ and indicates a
4 local MWP-1A amplitude of 12 to 22 m (ref. 1).

5 We applied the sea-level fingerprinting technique within a Bayesian statistical framework
6 to assess the likelihood of different MWP-1A source geometries (see Methods). Nine spatial
7 functions were defined to represent ice thickness changes across MWP-1A. Elastic-Earth sea-
8 level fingerprints were computed for each of these and then combined using different weighting
9 coefficients to test a large number of source scenarios (order 10,000) to satisfy statistical
10 requirements. The contribution of viscous Earth deformation due to ice-ocean loading and
11 rotational changes prior to MWP-1A was included via a model-correction to the observed MWP-
12 1A amplitudes described above (Supp. Table 1).

13 The nine spatial functions are based on deglaciation models of the Antarctic ice sheets
14 (AIS)^{21,22,23}, North American ice sheets (NAIS)^{12,24,25}, Fennoscandian ice sheet (FIS)²⁶, and
15 Greenland ice sheet (GIS)²⁷. Since the focus of this analysis is the AIS and NAIS, we
16 decomposed these ice complexes into several spatial functions, which are based on common
17 elements from different deglaciation models and so are relatively robust. The Antarctic
18 contribution to MWP-1A is defined using four spatial functions, corresponding to Wilkes Land,
19 the Weddell Sea, the Ross Sea, and the Antarctic Peninsula (Supp. Fig. 2). The North American
20 contribution is defined using three spatial functions based on recent modelling results for this ice
21 complex^{11,24,25} (Supplementary Fig. 3a-c). A single spatial function is defined for each of the FIS
22 and GIS, since the contribution of these ice sheets to MWP-1A was relatively minor and is less

debated¹¹. Their spatial functions are taken directly from recent reconstructions^{26,27} across the appropriate time window (Supplementary Fig. 3d,e).

For each spatial function except that of the NAIS saddle-collapse (Supp. Fig. 3c), the prior probability distribution of melt amplitude was taken as uniformly distributed between zero and twice the maximum melt contribution suggested in the source literature (Supplementary Table 3). For the saddle-collapse scenario, the upper bound of the amplitude prior was set equal to the estimated MWP-1A amplitude (15 m sea-level equivalent (sle))¹. The contributions from the AIS as a whole, the NAIS as a whole, and the FIS and GIS were treated as uncorrelated. Contributions from individual components of the AIS and of the NAIS were treated as uncorrelated prior to conditioning upon the total AIS or NAIS contribution.

We randomly sampled 40,000 individual MWP-1A source scenarios from the prior probability distribution (Supp. Table 3 and Supp. Figs 4 & 5). The likelihood of each scenario was then determined by comparing the calculated rsl rise to the model-corrected observations (Supp. Tables 1 & 2). Figure 3 shows the joint posterior probability distribution for the NAIS and AIS contributions when the original (Fig. 3a) and our revised (Fig. 3b) estimates of MWP-1A amplitude at Barbados and Sunda Shelf are adopted (the Tahiti amplitude range is the same in each case, 12-22.4 m (model corrected range from ref. 1)). As expected, the NAIS and AIS contributions are negatively correlated: as the contribution from one increases, less mass is required from the other. Posterior contribution estimates (95% probability) for the FIS and GIS are 0 to 2 m and 0 to 0.4 m sle, respectively (Table 1). These values are similar to the prior ranges (0-2.2 m and 0-0.4 m sle), indicating that the far-field data considered do not provide useful constraints on the contribution of these ice sheets to MWP-1A.

Two recent studies^{11,12} considered near-field evidence to constrain the NAIS MWP-1A contribution to either 2.8–3.7 m or 6.4–9.0 m sle (solid¹¹ and dashed-dotted¹² black boxes in Fig. 3a,c; the values provided in these studies, 6.7–8.7 over 800 years (ref. 11) and 9.4–13.2 m over 500 years (ref. 12), were scaled linearly to determine amplitudes for the 340-year interval from Tahiti¹). These estimates are based on both field and model constraints but apply different approaches to arrive at the ranges given. Rather than argue for the veracity of one over the other, we consider each to be equally plausible.

Jointly conditioning the prior probability distribution upon these alternative near-field constraints and the model-corrected MWP-1A original amplitudes inferred at Barbados² and Sunda Shelf³, as well as the more recent Tahiti constraint¹, indicates a 95% credible AIS contribution of either 5.9–10.1 m (ref. 11) or 2.1–9.1 m (ref. 12) sle (magenta curves in Fig. 3a), corresponding to a global mean sea-level rise of 11.2–16.1 m or 11.8–16.7 m, respectively. In comparison, using our revised far-field estimates leads to plausible MWP-1A source scenarios (Fig. 3b) with AIS contributions of 4.1–10.0 m (ref. 11) or 0–6.9 m (ref. 12) sle, and an estimated global mean sea-level rise of 9.3–14.6 m or 8.6–14.4 m sle, respectively (see Table 1 for a summary of results).

A recent fingerprinting analysis²⁸ to evaluate the plausibility of a large (~10 m) NAIS contribution to MWP-1A via a saddle collapse between the Cordilleran and Laurentide ice sheets demonstrated that the far-field constraints (different to those considered here for Barbados and Tahiti) are compatible with such a scenario. This study also concluded that the observations do not exclude the case of a dominant AIS contribution. The results in Fig. 3c are consistent with the results of ref. 28.

1 The 95% credible estimates of the local MWP-1A rsl amplitudes are 9.1-16.3 m at
2 Barbados, 11.7-17.9 m at Sunda Shelf, and 12.5-19.0 m at Tahiti (red and yellow ranges in Fig.
3 3c). The results in Fig. 3c indicate that the observed lower bound on MWP-1A amplitude at
4 Barbados and Tahiti and upper bound at Sunda Shelf provide the primary constraints on the
5 possible solution space. Therefore, new evidence from these locations that improve upon the
6 observational precision of these specific aspects of the local MWP-1A amplitude would reduce
7 the posterior uncertainties. As an example, increasing the value of the lower bound at Barbados
8 leads to estimates with a larger AIS contribution (Supp. Fig. 6). Note, however, that a relatively
9 large change in this value is required to markedly influence the results.

10 Our analysis conclusively demonstrates that, when data and model uncertainties are
11 carefully accounted for, the presently available far-field rsl reconstructions do not provide tightly
12 bounded constraints on MWP-1A partitioning: specifically, the 95% credible AIS contribution to
13 MWP-1A is 0–10.0 m sle when recent estimates of the NAIS contribution are considered^{11,12}.
14 Accordingly, our reassessment indicates that a significant AIS contribution may not be required,
15 thus potentially reconciling the apparent inconsistency between near-field¹⁰ and far-field
16 evidence. At the same time, however, our results suggest that a dominant AIS contribution
17 remains equally plausible. We note that any future improvements on the total NAIS contribution
18 can be directly applied to our AIS-NAIS partitioning diagram (Fig. 3c) and anticipate that the
19 approach taken here will provide the means to further constrain the source regions of MWP-1A
20 as more geological evidence becomes available. At present, uncertainty in the source distribution
21 of MWP-1A remains a primary limitation in our understanding of the causes and consequences
22 of this extreme event.

References

1. Deschamps, P. *et al.* Ice-sheet collapse and sea-level rise at the Bolling warming 14,600 years ago. *Nature* **483**, 559-564 (2012).
2. Peltier, W.R. & Fairbanks, R.G. Global glacial ice volume and Last Glacial Maximum duration from an extended Barbados sea level record. *Quat. Sci. Rev.* **25**, 3322-3337 (2006).
3. Hanebuth, T., Stattegger, K. & Grootes, P.M. Rapid flooding of the Sunda shelf: a late-glacial sea-level record. *Science* **288**, 1033-1035 (2000).
4. Golledge, N. R. *et al.* Antarctic contribution to meltwater pulse 1A from reduced Southern Ocean overturning. *Nat. Commun* **5**:5107 doi: 10.1038/ncomms6107 (2014).
5. Menviel, L., Timmermann, A., Timm, O. E. & Mouchet, A. Deconstructing the Last Glacial termination: the role of millennial and orbital-scale forcings. *Quat.Sci. Rev.* **30**, 1155–1172 (2011).
6. Stouffer, R. J., D. Seidov, D. & Haupt, B.J. Climate response to external sources of freshwater: North Atlantic versus the Southern Ocean, *J. Clim.* **20**(3), 436–448, doi:10.1175/JCLI4015.1 (2007).
7. Weaver, A.J., Saenko, O.A., Clark, P.U. & Mitrovica, J.X. Meltwater pulse 1A from Antarctica as a trigger of the Bølling-Allerød warm interval. *Science* **299**, 1709-1713 (2003).
8. Clark, P.U., Mitrovica, J.X., Milne, G.A. & Tamisiea, M.E. Sea-level fingerprinting as a direct test for the source of global meltwater pulse 1A. *Science* **295**, 2438-2441 (2002).
9. Bassett, S.E., Milne, G.A., Mitrovica, J.X. & Clark, P.U. Ice sheet and solid earth influences on far-field sea-level histories. *Science* **309**, 925-928 (2005).
10. Bentley, M.J. *et al.* A community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last Glacial Maximum. *Quat. Sci. Rev.* **100**, 1-9 (2014).
11. Carlson, A.E. & Clark, P.U. Ice sheet sources of sea level rise and freshwater discharge during the last deglaciation. *Rev. Geophys.* **50**, RG4007 (2012).
12. Tarasov, L., Dyke, A.S., Neal, R.M. & Peltier, W.R. A data-calibrated distribution of deglacial chronologies for the North American ice complex from glaciological modelling. *Earth Planet. Sci. Lett.* **315-316**, 30-40 (2012).
13. Mitrovica, J.X., Tamisiea, M.E., Davis, J.L., & Milne, G.A. Recent mass balance of polar ice sheets inferred from patterns of global sea-level change. *Nature* **409**(23), 1026-1029 (2001).
14. Fairbanks, R. G. A 17,000-year glacio-eustatic sea level record: Influence of glacial melting rates on the Younger Dryas event and deep-ocean circulation. *Nature* **342**, 637–642 (1989).
15. Bard, E., Hamelin, B. & Fairbanks, R.G. U-Th ages obtained by mass spectrometry in corals from Barbados: Sea level during the past 130,000 years. *Nature* **346**, 456–458 (1990).
16. Lighty, R.G. Fleishy-algal domination of a modern Bahamian barrier reef: example of an alternate climate reef community. *Proc. 4th Int. Coral Reef Symp. Manila* **1**, 772 (1981).
17. Neumann, A.C., & McIntyre, I.G. Reef response to sea-level rise: Keep-up, catch-up, or give-up. *Proc. 5th Int. Coral Reef Cong. Tahiti* **3**, 105-110 (1985).

18. Lambeck, K., Yokoyama, Y. & Purcell, T. Into and out of the Last Glacial Maximum: sea-level change during Oxygen Isotope Stages 3 and 2. *Quat. Sci. Rev.* **21**, 343-360 (2002).
19. Bard, E., Hamelin, B., Arnold, M., Montaggioni, L., Cabioch, G., Faure, G. & Rougerie, F. Deglacial sea-level record from Tahiti corals and the timing of global meltwater discharge, *Nature* **382**, 241–244 (1996).
20. Blanchon, P. & Blakeway, D. Are catch-up reefs and artifact of coring? *Sedimentology* **50**, 1271-1282 (2003).
21. Briggs, R. D., Pollard, D. & Tarasov, L. A data-constrained large ensemble analysis of Antarctic evolution since the Eemian, *Quat. Sci. Rev.* **103**, 91–115 (2014).
22. Gomez, N., Pollard D. & Mitrovica, J.X. 2013. A 3-D coupled ice sheet-sea level model applied to Antarctica through the last 40 ky. *Earth Planet. Sci. Lett.* **384**, 88-99 (2013).
23. Whitehouse, P.L., Bentley, M.J. & Le Brocq, A.M. A deglacial model for Antarctica: geological constraints and glaciological modelling as a basis for a new model of Antarctic glacial isostatic adjustment. *Quat. Sci. Rev.* **32**, 1-24 (2012).
24. Carlson, A.E. *et al.* Modeling the surface mass-balance response of the Laurentide Ice Sheet to Bølling warming and its contribution to Meltwater Pulse 1A. *Earth Planet. Sci. Lett.* **315-316**, 24-29 (2012).
25. Gregoire, L.J., Payne, A.J. & Valdes, P.J. Deglacial rapid sea-level rises caused by ice-sheet saddle collapses. *Nature* **487**, 219-222 (2012).
26. Lambeck, K. Smither, C. & Johnston, P. Sea-level change, glacial rebound and mantle viscosity for northern Europe. *Geophys. J. Int.* **134**, 102-144 (1998).
27. Simpson, M.J.R., Milne, G.A., Huybrechts, P. & Long, A.J. Calibrating a glaciological model of the Greenland ice sheet from the Last Glacial Maximum to present-day using field observations of relative sea level and ice extent. *Quat. Sci. Rev.* **28**, 1631-1657 (2009).
28. Gomez, N., Gregoire, L. J., Mitrovica, J. X. & Payne, A. J. Laurentide-Cordilleran Ice Sheet saddle collapse as a contribution to meltwater pulse 1A, *Geophys. Res. Lett.*, **42**, 3954–3962, doi:10.1002/2015GL063960 (2015).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgements

J.L. and G.A.M. are supported by the Natural Sciences and Engineering Research Council of Canada. G.A.M. is also supported by the Canada Research Chairs program. R.E.K. acknowledges support from the National Science Foundation (NSF) of the United States (ARC 1203415). P.U.C. acknowledges support from the NSF Antarctic Glaciology Program (ANT1260719) and the NSF Marine Geology and Geophysics Program (OCE1335197). We thank Jerry Mitrovica for providing information that contributed to the work presented in this paper. This work benefitted from discussions at several PALeo constraints on SEA level rise (PALSEA) workshops.

Author Contributions

J.L. performed the research and led the writing of the manuscript. G.A.M., R.E.K, P.U.C., and I.S. advised J.L. in performing the research and contributed to the writing of the manuscript.

Author Information

Reprints and permissions information is available at www.nature.com/reprints. Correspondence and requests for materials should be addressed to gamilne@uottawa.ca.

Figure and Table Legends

Figure 1 Illustration of method used to estimate MWP-1A amplitude at Barbados using age and depth information from coral samples. Horizontal bars denote age uncertainties and hard lower bounds. Vertical bars denote depth uncertainties. Purple points are from site 9, orange from Site 12, and black from site 15 of Peltier and Fairbanks². Grey box denotes MWP-1A timing based on the Tahitian record of Deschamps et al.¹. A maximum MWP-1A sea-level change is set by lowest slope that is consistent with the observations within uncertainty of sea level after MWP-1A (blue dotted line), and a minimum sea-level change is set by the steepest consistent slope (red dotted line). Since the first two sample observations plotted (black and left-most purple index points) are the same age within uncertainty, we took the overlapping depth range and total combined age range for these two index points to define our earliest data constraint and extrapolated back to 14.65 ka, the earliest that MWP-1A could have begun¹, to get the starting depth of MWP-1A. The solid blue and red lines show how the MWP-1A amplitudes were determined from extrapolation. Thick blue and red bars denote the corresponding estimates of maximum and minimum MWP-1A amplitudes. The solid green line indicates the result of Deschamps et al.¹, who did not consider data uncertainty and extrapolated back in time to the first index point shown, rather than 14.65 ka as we have done. The thick green bar shows the MWP-1A amplitude estimated by Deschamps et al.¹.

Figure 2 Relative sea level reconstructions for the Sunda Shelf and model estimate of sea-level gradient across this region at the time of MWP-1A. (a) Sea-level contour lines across Sunda Shelf are the mean of a model ensemble (see Methods). Black star denotes site 18300, to which all other sites are reduced; cyan dot marks site 18299, blue 18301, magenta 18302, purple 18307, red 18308, green 18309, and tan 18310 from the Hanebuth et al.³ study. (b) and (c) show relative sea-level constraints at Sunda Shelf before and after the spatial correction is made, with colours corresponding to different core sites as defined in (a) and yellow stars marking *in-situ* samples. Horizontal bars mark age uncertainty. Vertical bars mark depth uncertainty, which includes GIA model uncertainty in (c). Blue and red bars depict the maximum and minimum local MWP-1A amplitude. Grey box denotes MWP-1A timing¹.

Figure 3 Posterior distribution of NAIS and AIS sea-level contributions conditioned on far-field rsl reconstructions. Results for previously-published far-field MWP-1A amplitude estimates^{1,2,3} (a) and for our revised amplitude estimates (c), with solution density indicated by the color scale. The magenta contour indicates the central 95% credible range. The black

outlines indicate two recent estimates of the NAIS contribution to MWP-1A based on near-field evidence: 2.8-3.7 m sle (solid line; ref. 11) and 6.4-9.0 m sle (dashed-dotted line; ref. 12). In (b) and (d), the thin vertical bars denote the MWP-1A amplitudes and uncertainties at each of far-field sites corresponding to (a) and (c), respectively, while the colored bars show the local MWP-1A amplitudes produced by scenarios that satisfy all far-field constraints. Cyan, yellow, and red bars show the 99%, 95%, and 67% credible intervals, respectively. Note that the model-corrected upper bound of MWP-1A amplitude at Barbados (33.6 m) is not visible.

Table 1 Posterior estimates on MWP-1A source partitioning. Quoted results are 95% credible intervals. Ice volumes given as metres of sea-level equivalent calculated using the present-day ocean area.

Methods

Sea-Level Model

The glacial isostatic adjustment (GIA) model adopted in this study computes sea-level changes due to solid Earth deformation and gravity changes associated with the redistribution of ice and water on the Earth's surface²⁹⁻³¹. In addition to the ice-ocean loading, the model also includes the influence of changes in Earth rotation due to GIA³²⁻³⁴ as this can contribute significantly to the sea-level response, particularly during rapid and large events such as Meltwater Pulse 1A (MWP-1A)^{8,28,35}.

The two primary inputs to the model are a space-time reconstruction of grounded ice thickness and a model of Earth sub-surface density and rheology structure. Different ice models are applied in this study and they are defined and described where appropriate. The adopted Earth model is spherically symmetric and so includes only changes in parameters with depth. The elastic and density depth profiles are taken from a seismic model³⁶ and are defined with a depth resolution of 5-25 km. These profiles were not varied in this analysis. The viscous structure is less precisely known and so a large range of parameters were considered (details below where appropriate). Given the relatively large uncertainty in this model aspect, the depth parameterisation of the viscosity profile was considerably lower resolution compared to that for the elastic and density changes. Following a number of previous GIA analyses, we define an outer shell with very high viscosity (10^{43} Pas) to simulate an elastic lithosphere; the thickness of this outer shell is varied in the modelling. We define an "upper mantle" region from the base of the model lithosphere to 670 km depth and a "lower mantle" region from 670 km to the Core-Mantle boundary. Viscosity is defined to be uniform in these two regions.

Determining MWP-1A Amplitude From the Sunda Shelf Sea-Level Reconstructions

Given the relatively large spatial spread in the locations where relative sea level (rsl) was reconstructed on the Sunda Shelf, it is necessary to reduce the observations to a single locality in order to accurately determine the local MWP-1A amplitude. We applied the model described in Section 1 for this purpose and computed rsl in the region for a total of 324 parameter sets

1 comprising two ice models (ICE5G³⁷ and that of Bassett et al⁹) and 162 Earth viscosity models
2 (lithosphere thickness of 71, 96, and 120 km; upper mantle viscosity of 0.1, 0.2, 0.3, 0.5, 0.8, and
3 1×10^{21} Pas; and lower mantle viscosity of 1, 2, 3, 5, 8, 10, 20, 30, and 50×10^{21} Pas). We used
4 the mean difference between the sea-level at each core location and that at core site 18300 to
5 define a spatial correction for each rsl data point, calibrated with IntCal13³⁸, around the MWP-
6 1A period (Fig 2a, main text). The uncertainty in the model correction was taken to be the spread
7 in results produced by the parameter ranges defined above. By reducing the rsl index points to a
8 single location (core site 18300: 4.3630° N, 108.6536° E), we found a revised MWP-1A
9 amplitude of 7.5 to 17.3 m, compared with 12 to 20 m for the uncorrected data.

10
11 We are confident that the range in model parameters considered provides a conservative estimate
12 of the model uncertainty in the spatial correction applied. Given that the primary contributor to
13 the spatial rsl gradient in this region is ocean loading, the sensitivity of the results to the ice
14 history is largely through the time variation in the global ice volume (or sea level equivalent, sle)
15 rather than differences in the spatial distribution of ice through time. Both of the ice models
16 adopted have been calibrated to fit far-field RSL observations for considerably different Earth
17 viscosity models, leading to significant differences in their respective sle curves^{9,37}. Furthermore,
18 the two models are based on contrasting source scenarios for MWP-1A (one³⁷ solely northern
19 and the other⁹ dominantly southern). Therefore, we believe that these two models likely bound
20 the uncertainty associated with the aspect of the ice model that influences the modelled ocean
21 loading. With regard to the Earth model viscosity structure, the parameter ranges adopted likely
22 overestimate the uncertainty in this model input.

23 24 25 **Contribution of viscous Earth deformation to the sea-level fingerprints**

26
27 Spatial patterns of rsl change associated with melting bodies of land ice are governed by the
28 geographic distribution of ice and the associated deformational response of the solid Earth³⁹.
29 Over relatively short timescales (a few centuries), the contribution of viscous Earth deformation
30 to the pattern of rsl change is relatively small compared to changes over longer time periods
31 (multi-millennial to deglacial) that have been more commonly considered in GIA modelling
32 studies. Thus, a primary benefit of short-time-scale problems such as MWP-1A is that sensitivity
33 to Earth viscosity is relatively low^{8,28} and so the considerable uncertainty in this model parameter
34 is less influential on the results. However, viscous deformation can contribute as much as a metre
35 or so to the computed sea-level fingerprints^{8,28} and so we consider its impact by estimating and
36 then removing it from the far-field rsl constraints of local MWP-1A amplitude.

37
38 There are two components of viscous solid Earth deformation that contribute to the spatial
39 pattern (or fingerprint) of rsl change during MWP-1A: that associated with ice-ocean loading
40 and rotational changes prior to the event and that due to these changes during the event. We
41 consider only the former as computing the viscous deformation associated with the large number
42 of source scenarios (10s of thousands) required to ensure our results were statistically robust is
43 computationally prohibitive.

44
45 The magnitude of the pre-MWP-1A viscous “overprint” depends on a number of factors,
46 including the amplitude and timing of the loading and rotational changes prior to MWP-1A, the

viscosity structure of the Earth and the duration of MWP-1A²⁸ (the longer the duration, the larger the viscous contribution will be). We computed the viscous response due to loading before MWP-1A at all three sites using the suite of ice and Earth model parameters described in Section 2 (324 model runs in total) by running the full time history of the ice model: from the end of the last interglacial up to 14.5 ka. The model was then run for an additional time step of 500 years with no further loading or rotational changes to determine the viscous contribution over the period 14.5 to 14.0 ka. Given that the viscous signal is approximately linear over this period²⁸, we scaled the results to be representative of a 340 year interval as adopted elsewhere in this analysis. Our results (Supplementary Fig. 1), agree in sign and are similar in amplitude to those in ref. 28 (see their Fig. 3). However, since we neglected the viscous deformation during MWP-1A, the mean of our model spread is less than the values presented in ref. 28.

The pre-MWP-1A viscous signal shown in Supplementary Fig. 1 was incorporated into our final results (Fig. 3, main text) by considering the full range of the model spread. The model spread was combined directly to the observed values in order to produce a conservative estimate of the uncertainty associated with the pre-MWP-1A viscous contribution. The raw and model-corrected MWP-1A amplitudes are given in Supplementary Table 1. To test the impact of pre-MWP-1A viscous deformation on our final results, we computed the posterior probability estimates without applying this model correction (i.e. ignoring all viscous effects) (Supplementary Table 2). The results show that the estimated AIS contribution is affected but those for the FIS and GIS are not. The differences in the AIS 95%-credible ranges, with and without the viscous correction, are relatively small and depend on the adopted range for the NAIS contribution.

Melt source geometries

To compute rsl fingerprints associated with ice sheet changes during MWP-1A, it is necessary to define the melt source geometries to be tested. As described in the main text, we did this by specifying nine spatial functions identified from a number of recent ice model reconstructions. For Antarctica and North America, specific source regions within these ice complexes were defined (Supplementary Figs 2 & 3a-c, respectively). For these regions, more than one model reconstruction was considered (see main text) so as to determine source regions that are compatible with multiple studies and thus more robust. In contrast, the melt distributions for Fennoscandia and Greenland were taken directly from single studies (see main text; Supplementary Fig. 3d,e) given that their contribution to MWP-1A is relatively minor and less debated¹¹.

We note that the spatial functions defined in Supplementary Figs 2 & 3 have relatively crude spatial fidelity as they were not intended to accurately define the changes in ice distribution during MWP-1A. Rather, they were intended only to provide an approximate representation of these changes for each region. While our final results (Fig. 3, main text) indicate that the far-field rsl constraints show a clear sensitivity to the partitioning of mass loss between the Antarctic and North American ice sheets, their sensitivity to the partitioning of mass loss within these regions is much less pronounced, particularly for Antarctica. Therefore, we believe that the spatial fidelity of the AIS and NAIS source functions is more than adequate given the limited geographic distribution and precision of the rsl data considered.

The nine functions defining ice changes during MWP-1A were used as input to the GIA sea-level model to compute the rsl rise at each of the three far-field sites for the case of an elastic Earth rheology. The computed rise at each site was normalised by the volume of ice loss (in metres sle) to define a “fingerprint” for each melt source.

Statistical Methodology

We quantify the Bayesian probability of different alternative source region contributions to MWP-1A. If \mathbf{H} is a particular set of ice sheet contributions, \mathbf{F} the far-field observational constraints, and \mathbf{N} the near-field observational constraints, then by Bayes’ theorem,

$$P(\mathbf{H}|\mathbf{F},\mathbf{N}) \sim P(\mathbf{F},\mathbf{N}|\mathbf{H}) P(\mathbf{H}) \quad (1)$$

To estimate the posterior probability distribution $P(\mathbf{H}|\mathbf{F},\mathbf{N})$, we took 40,000 maximin Latin hypercube samples from the prior probability distribution $P(\mathbf{H})$, which is described below, and weight each sample by its likelihood, $P(\mathbf{F},\mathbf{N}|\mathbf{H})$. We assume that the far-field observations have uniform likelihoods in terms of rsl (which is a linear transformation of \mathbf{H} , generated using the spatial functions described above). In particular, we assume that Barbados, Sunda Shelf and Tahiti have likelihoods that are, respectively, uniform between 9.0-33.6 m, 7.5–17.9 m, and 12.0–22.4 m rsl (Supplementary Table 1). We further assume that the near-field observations have uniform likelihoods in terms of ice volume; thus, they serve simply to truncate the posterior distribution calculated by conditioning on far-field distributions. As a result of the uniform likelihoods, each sample from $P(\mathbf{H})$ has a relative weight of either zero or $1/n$, where n is the total number of samples with non-zero likelihoods.

The priors for the individual source regions are shown in Supplementary Table 3. To help account for differences in the interpretation of near-field data¹¹ and to remain consistent with the conservative nature of this analysis, the upper bound to the uniform prior for eight of the nine source regions was set equal to twice that indicated in the source literature. For the region that represents the saddle collapse signal^{12,25} (Supplementary Fig. 3c), the upper bound for the uniform prior was set equal to 15 m sle (Supplementary Table 3). For each component source region in the AIS and NAIS, we used a uniform prior that is conditioned upon the uniform prior for the ice sheet as a whole; these were sampled by first sampling from the prior for the ice sheet as a whole, then randomly dividing the ice sheet into sections and rejecting those divisions incompatible with the uniform priors for the individual source regions.

Supplementary Fig. 4 shows the sampling density for the NAIS versus the AIS as well as histograms indicating the number of samples for a given total contribution from each of the four source regions. From the sampled total contribution of the AIS and NAIS, contributions from the sub-sectors were sampled until all sub-sector constraints compatible with the specified total AIS/NAIS contribution were satisfied. Supplementary Fig. 5 provides histograms of the number of times a given sub-sector contribution was sampled.

Code availability: The code used for the statistical analysis is available upon request.

References

- 1
2
3 29. Farrell, W.E. & Clark, J.A. On postglacial sea-level. *Geophys. J. R. Astr. Soc.* 46, 647–667
4 (1976).
5 30. Mitrovica, J.X. & Milne, G.A. On post-glacial sea level - I. General theory. *Geophys. J. Int.*
6 154, 253–267 (2003).
7 31. Kendall, R., Mitrovica, J.X. & Milne, G.A. On post-glacial sea level — II. Numerical
8 formulation and comparative results on spherically symmetric models. *Geophys. J. Int.* 161,
9 679–706 (2005).
10 32. Sabadini, R., Yuen, D.A. & Boschi, E. Polar wander and the forced responses of a rotating,
11 multi-layered, viscoelastic planet. *J. Geophys. Res.* 87, 2885–2903 (1982).
12 33. Milne, G.A. & Mitrovica, J.X. Postglacial sea-level change on a rotating Earth. *Geophys. J.*
13 *Int.* 133, 1–19 (1998).
14 34. Mitrovica, J.X., Wahr, J., Matsuyama, I. & Paulson, A. The rotational stability of an ice-age
15 earth. *Geophys. J. Int.* 161, 491–506 (2005).
16 35. Mitrovica, J.X., Gomez, N., Morrow, E., Hay, C., Latychev, K. & Tamisiea, M.E. On the
17 robustness of predictions of sea level fingerprints. *Geophys. J. Int.* 187, 729–742 (2011).
18 36. Dziewonski, A.M. & Anderson, D.L. Preliminary reference earth model (PREM). *Phys.*
19 *Earth Planet. Inter.* 25, 297–356 (1981).
20 37. Peltier, W.R. Global glacial isostasy and the surface of the ice-age Earth: the ice-5G (VM2)
21 model and GRACE. *Annu. Rev. Earth Planet Sci.* 32, 111–149 (2004).
22 38. Reimer, P.J. et al. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years
23 cal BP. *Radiocarbon* 55(4), 1869–1887 (2013).
24 39. Clark, J.A., Farrell, W.E. & Peltier, W.R. Global changes in postglacial sea level: a
25 numerical calculation. *Quat. Res.* 9, 265–287 (1978).
26

Figure 1




Figure 2





Figure 3



Sea-Level Constraints on the Amplitude and Source Distribution of Meltwater Pulse

1A: Supplementary Information

Jean Liu¹, Glenn A. Milne^{1*}, Robert E. Kopp², Peter U. Clark³, and Ian Shennan⁴

¹Department of Earth and Environmental Sciences, University of Ottawa

²Department of Earth & Planetary Sciences, Rutgers Energy Institute, and Institute of Earth,
Ocean & Atmospheric Sciences, Rutgers University

³ College of Earth, Ocean, and Atmospheric Sciences, Oregon State University

⁴Department of Geography, Durham University

Location	Observed MWP-1A amplitude (m)	Model-corrected MWP- 1A amplitude (m)
Barbados	9.7 to 33.6	9.0 to 33.6
Sunda Shelf	7.5 to 17.3	7.5 to 17.9
Tahiti	12.0 to 22.0	12.0 to 22.4


Supplementary Table 1: MWP-1A amplitude estimates with and without model-correction for solid Earth viscous contribution associated with ice-ocean loading and rotational changes prior to MWP-1A. Note that ranges given include the model spread shown in Supplementary Fig 1. The model-corrected values were used to generate the results in Fig. 3 (main text).

Ice Sheet(s)	Prior Distribution	Given NAIS constraints of 2.8-3.7 m (ref. 11)		Given NAIS constraints of 6.4-9.0 m (ref. 12)	
		Including Viscous Signal (as in main text)	Not Including Viscous Signal	Including Viscous Signal (as in main text)	Not Including Viscous Signal
AIS	0 to 10.2 m	4.1 to 10.0 m	3.5 to 10.1 m	0 to 6.9 m	0 to 6.5 m
NAIS	0 to 32.4 m	2.8 to 3.7 m	2.8 to 3.7 m	6.4 to 9.0 m	6.4 to 9 m
FIS	0 to 2.2 m	0 to 2.2 m	0 to 2.2 m	0 to 2.2 m	0 to 2.2 m
GIS	0 to 0.4 m	0 to 0.4 m	0 to 0.4 m	0 to 0.4 m	0 to 0.4 m
TOTAL	0 to 45.2 m	9.3 to 14.6 m	8.9 to 14.0 m	8.6 to 14.4 m	9.1 to 13.9 m


Supplementary Table 2: Constraints on MWP-1A source partitioning with and without incorporating model estimates of the pre-MWP-1A viscous contribution (see Supplementary Table 1). Ice volumes are given as metres of sea-level equivalent calculated using the present-day ocean area.

Ice Sheet	Spatial Weighting Function	Corresponding Figure	Prior Probability Distribution (m)
AIS	Wilkes Land	Supp. Fig. 2a	U(0, 2.0)
	Weddell	Supp. Fig. 2b	U(0, 2.4)
	Ross	Supp. Fig. 2c	U(0, 3.0)
	Peninsula	Supp. Fig. 2d	U(0, 2.8)
	Total AIS		U(0, 10.2)
NAIS	Localized signal in Northwest	Supp. Fig. 3a	U(0, 3.8)
	Broader signal of regional mass losses and gains	Supp. Fig. 3b	U(0,13.6)
	“Saddle collapse” separating LIS and CIS	Supp. Fig. 3c	U(0,15.0)
	Total NAIS		U(0,32.4)
FIS	Full signal	Supp. Fig. 3d	U(0, 2.2)
GIS	Full signal	Supp. Fig. 3e	U(0, 0.4)


Supplementary Table 3: Prior melt volume ranges (in sea-level equivalent) for each of the spatial area weighting functions considered. Ice volumes are given as metres of sea-level equivalent calculated using the present-day ocean area.




Supplementary Fig. 1: Three hundred and twenty four model realisations of the viscous sea-level response across the MWP-1A time window due to ice-ocean loading and rotational changes prior to this event at Barbados **(a)**, Sunda Shelf **(b)** and Tahiti **(c)**. Model parameter values are described in Methods.




Supplementary Fig. 2: Antarctic spatial melting functions based on recent models of AIS deglaciation^{21,22,23}. These define the spatial distribution of ice melt for: **(a)** Wilkes Land, **(b)** Weddell Sea, **(c)** Ross Sea, and **(d)** Antarctic Peninsula. Note that the functions in (c) and (d) are spatially uniform within the areas indicated. The maximum volume loss for each sector is indicated in Supplementary Table 3.




Supplementary Fig. 3: Northern Hemisphere spatial melting functions. These functions define the modelled melt distribution from: (a) the northwest of the NAIS¹², (b) the broader mass loss and gains across the NAIS²⁴, (c) the saddle collapse separation of the CIS and LIS^{12,25} (d) the FIS²⁶, and (e) the GIS²⁷. Note that the functions in (a) and (c) are spatially uniform within the areas indicated. The maximum volume loss for each sector is indicated in Supplementary Table 3.



Supplementary Fig. 4: Sampling distribution from the prior probability. Latin Hypercube sampling of North American vs. Antarctic contributions (left) and from all source regions considered (right).



Supplementary Fig. 5: Sampling distribution of sub-sectors of the AIS (top) and NAIS (bottom).



Supplementary Fig. 6: Posterior estimates (95% and higher probability) for different values of the minimum MWP-1A amplitude at Barbados (as indicated by the different colours). The black outlines indicate two recent estimates of the NAIS contribution to MWP-1A based on near-field evidence: 2.8-3.7 m sle (solid line; ref. 11) and 6.4-9.0 m sle (dashed-dotted line; ref. 12)